

Crescentic glomerulonephritis in a man with a history of methamphetamine abuse; a possible cause for concern



Dorsa Jahangiri¹ , Mohammadreza Ardalan^{2*} , Muhammed Mubarak³ , Shahrzad Alimohammadi^{4,5} ,
Hamid Reza Jahantigh^{6,7} , Sanam Saeifar⁸ 

¹Independent Researcher, 43185 Cardston Place Leesburg Virginia, 20176, USA

²Kidney Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

³JIK Department of Histopathology, Sindh Institute of Urology and Transplantation, Karachi, Pakistan

⁴Doctoral School of Molecular Medicine, University of Debrecen, Debrecen, Hungary

⁵Department of Immunology, Faculty of Medicine, University of Debrecen, Debrecen, Hungary

⁶Interdisciplinary Department of Medicine - Section of Occupational Medicine, University of Bari, Bari, Italy

⁷Animal Health and Zoonosis PhD Course, Department of Veterinary Medicine, University of Bari, Bari, Italy

⁸Buchmann Institute for Molecular Life Sciences (BMLS), Cluster of Excellence Frankfurt Macromolecular Complexes (CEF-MC), Goethe University Frankfurt am Main, Frankfurt am Main, Germany

ARTICLE INFO

Article Type:

Photonephrology

Article History:

Received: 2 July 2022

Accepted: 18 September 2022

Published online: 8 October 2022

Keywords:

Crescentic, Methamphetamine
Acute kidney injury

Implication for health policy/practice/research/medical education:

Methamphetamine abuse typically causes acute kidney injury mainly due to rhabdomyolysis, malignant hypertension and volume depletion and acute tubular necrosis. However, methamphetamine-induced crescentic glomerulonephritis (CresGN) has not been reported so far.

Please cite this paper as: Jahangiri D, Ardalan M, Mubarak M, Alimohammadi S, Jahantigh HR, Saeifar S. Crescentic glomerulonephritis in a man with a history of methamphetamine abuse; a possible cause for concern. J Nephropharmacol. 2023;12(1):e10508. DOI: 10.34172/npj.2022.10508.

A 48-year-old man was referred to our nephrology clinic due to hematuria, proteinuria, coupled with elevated serum creatinine level (6.8 mg/dL). He reported a recent history of methamphetamine abuse by smoke inhalation. When admitted, his blood pressure was 150/90 mm Hg, and the report of fundoscopic examination was normal. Further laboratory examinations revealed: white blood cells (WBC) count, $8700 \times 10^3/\text{mm}^3$, platelets, $186 \times 10^3/\text{mm}^3$, hemoglobin; 9.6 g/dL, urea 145 mg/dL, sodium, 137 mEq/L, potassium, 4.7 mEq/L, calcium, 9 mg/dL, and phosphorus, 5.5 mg/dL. Antinuclear antibody (ANA), anti-double-stranded DNA (anti-dsDNA), and anti-neutrophil cytoplasmic antibody (c-ANCA and p-ANCA) were all negative. The complement CH50 level was low; 15.4 U/ml (Normal range: 51-150 U/ml), but both C3 and C4 levels were in the normal range. The serological tests for viral hepatitis B, C, and human immunodeficiency virus (HIV) were all negative.

An ultrasound-guided renal biopsy was conducted. Light microscopy revealed cellular crescents in six

glomeruli and fibrous crescent in one among a total of 12 sampled glomeruli (Figure 1). Immunofluorescence microscopy showed mild capillary wall staining for IgG (+) and C3 (+ +), but negative for C1q, IgM, IgA, kappa, and lambda. The patient received pulse therapy with 500 mg/day methylprednisolone for three days followed by oral prednisone and cyclophosphamide intravenous (IV) pulse. Three weeks later, his serum creatinine level decreased to 2.1 mg/dL, the CH50 level returned to its normal range and his general condition improved significantly.

Methamphetamine abuse typically causes acute kidney injury mainly due to rhabdomyolysis, malignant hypertension and volume depletion and acute tubular necrosis. However, methamphetamine-induced crescentic glomerulonephritis (CresGN) has not been reported so far. Unfortunately, establishing the causal relationship of CresGN to methamphetamine nephrotoxicity in his case was impossible. Although, it illustrates the necessity of considering this drug as a possible cause of




Figure 1. Two glomeruli show prominent cellular crescents formation compressing the native glomerular tufts. (A) The crescent partially covers the tuft. There is mild interstitial inflammatory cell infiltrate in the surrounding parenchyma (Jones' silver stain $\times 200$). (B) The crescent is circumferentially covering the entire tuft. Mild periglomerular inflammatory infiltrate is present (Jones' silver stain $\times 400$).

renal damage in such situations. Whilst it is difficult to prove that methamphetamine may be adulterated with other compounds such as cocaine and levamisole, that are known for their nephrotoxic pauci-immune CresGN causation (1-3).

Authors' contribution

Conceptualization: MA, Methodology: MM, ShA & HRJ, Validation: DJ, Formal Analysis: MM, Investigation: MA, Resources: DJ & SS, Data Curation: MA, Writing—Original Draft Preparation: MA & SS, Writing—Review and Editing: DJ & SS, Visualization: DJ & SS, Supervision: MA, Project Administration: MA, Funding Acquisition: -

Conflicts of interest

The authors declare that they have no conflicts of interest.

Ethical issues

This photonephrology was conducted in accord with the World Medical Association Declaration of Helsinki. The patient has given us a written informed consent for publication as a photonephrology. Besides, ethical issues

(including plagiarism, data fabrication, and double publication) have been completely observed by the authors.

Funding/Support

None.

References

1. Pendergraft WF 3rd, Herlitz LC, Thornley-Brown D, Rosner M, Niles JL. Nephrotoxic effects of common and emerging drugs of abuse. *Clin J Am Soc Nephrol.* 2014;9:1996-2005. doi: 10.2215/CJN.00360114.
2. Carrara C, Emili S, Lin M, Alpers CE. Necrotizing and crescentic glomerulonephritis with membranous nephropathy in a patient exposed to levamisole-adulterated cocaine. *Clin Kidney J.* 2016;9:234-8. doi: 10.1093/ckj/sfv141.
3. Baradhi KM, Pathireddy S, Bose S, Aeddula NR. Methamphetamine (N-methylamphetamine)-induced renal disease: underevaluated cause of end-stage renal disease (ESRD). *BMJ Case Rep.* 2019;12:e230288. doi: 10.1136/bcr-2019-230288.

Copyright © 2023 The Author(s); Published by Society of Diabetic Nephropathy Prevention. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.